



King's Research Portal

DOI:

[10.1186/2046-7648-4-S1-A121](https://doi.org/10.1186/2046-7648-4-S1-A121)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

O'Brien, K. A., Pollock, R., Stroud, M., Kumar, A., Lambert, R. J., Green, D. A., Edwards, L. M., & Harridge, S. (2015). Physiological and metabolic responses to prolonged hypoxia and extreme cold: Preliminary data from the White Mars Antarctica winter expedition. *Extreme physiology & medicine*, 4(1). <https://doi.org/10.1186/2046-7648-4-S1-A121>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

MEETING ABSTRACT

Open Access

Physiological and metabolic responses to prolonged hypoxia and extreme cold: Preliminary data from the White Mars Antarctica winter expedition

Katie A O'Brien^{1*}, Ross Pollock¹, Mike Stroud², Alex Kumar³, Robert J Lambert⁴, David A Green¹, Lindsay M Edwards¹, Stephen Harridge¹

From 15th International Conference on Environmental Ergonomics (ICEE XV)
Portsmouth, UK. 28 June - 3 July 2015

Introduction

The Antarctic winter is amongst the most extreme environments on earth. Human adaptation to this environment, where severe cold is coupled with moderate altitudes, is poorly understood. In this study, a number of physiological and metabolic measurements were made on a small group of trekkers before and after an attempted winter crossing of Antarctica (White Mars Expedition).


Methods

5 male subjects aged 28-54 yrs were assessed prior to and following a 24 week stay in Antarctica, including 14 weeks above 2,500 m. Measurements included

assessment of body fat and bone mineral density (DXA), cardiorespiratory responses to an incremental exercise test, lung and cardiovascular function as well as metabolic analysis of serum using ¹H-NMR spectroscopy.

Results

Significant changes were found in the following parameters pre to post expedition, identified using a paired Student t test (mean (SD), $p < 0.05$). There was an increase in % lean tissue (79+4 vs. 81+3%), a decrease in % fat tissue (21(4) vs. 19(3) %) and body fat mass (16(5) vs. 14(4) kg), although whole body weight did not change. Both spine bone mineral density (1.2(0.05) vs. 1.13(0.04) g.cm²) and FEV₁:FVC (68(10) vs. 62(8))



* Correspondence: katie.obrien@kcl.ac.uk

¹Centre of Human and Aerospace Physiological Sciences, King's College London, London, UK

Full list of author information is available at the end of the article

were decreased. $\text{VO}_{2\text{max}}$ did not significantly change from the pre-expedition 42mL.kg.min, however an increase was observed pre to post expedition in the respiratory exchange ratio (RER) at each stage (10%) of the $\text{VO}_{2\text{max}}$ test (Figure 1A). Metabolomics analysis of serum samples revealed changes in two peaks within principal component 2: glucose and a fatty acid CH_2 resonance (Figure 1B and 1C).

Discussion

These results are suggestive of a number of physiological changes resulting from prolonged exposure to the Antarctic winter. In particular, we observed a change in metabolic signature involving changes to both glucose and fatty acid homeostasis with a shift towards increased reliance on carbohydrate metabolism during exercise.

Conclusion

This study has highlighted areas of interest for future investigations into the physiological responses to this unique environment.

Authors' details

¹Centre of Human and Aerospace Physiological Sciences, King's College London, London, UK. ²NIHR Biomedical Research Centre for Nutrition, Southampton, UK. ³University Hospitals of Leicester, UK & University of Fribourg, Switzerland. ⁴Dept. of Trauma and Orthopaedic Surgery, Royal Infirmary of Edinburgh, UK.

Published: 14 September 2015

doi:10.1186/2046-7648-4-S1-A121

Cite this article as: O'Brien *et al.*: Physiological and metabolic responses to prolonged hypoxia and extreme cold: Preliminary data from the White Mars Antarctica winter expedition. *Extreme Physiology & Medicine* 2015 **4**(Suppl 1):A121.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

